SPREAD OUT TO STOP THE SPREAD

- Practice social distancing at all times and stay six feet away from your colleagues.
- Wear a surgical mask at all times, unless in a room alone.
- In the reception, pharmacy, lab or radiology areas, use a physical marker, such as an "X" on the floor, to indicate a six-foot distance from where the patients should wait.

Additional protective measures:

- The majority of patients are using virtual visits. Patients will only be seen in the clinics if
 it is medically necessary.
- Ensure that signs are posted discouraging patients with respiratory symptoms from entering the clinic.
- Front desk staff should wear surgical masks if receiving walk-ins.
- Walk-ins with respiratory symptoms should immediately be given a mask and placed in a separate room.
- Frequently disinfect areas that patients and staff have contact with, including door handles, tablets, keyboards, exam beds, multi-use devices, desk and cell phones etc.
- Practice hand hygiene often; COVID-19 is destroyed by proper hand hygiene.
- Avoid touching your face, especially your mouth and mucous membranes.
- Cover your cough.
- ALL direct patient care must be conducted wearing an N95 mask.
- Do not touch the inside of masks; reuse until visibly soiled or structural integrity lost.
- covidence of the control of the cont
- Team member temperatures are taken daily upon arrival to work, and anyone with a
 fever 100.0 degrees or greater will be sent home.